

Impacts of Regional Land Use and Land Cover on Rainfall: An Overview

R.A. PIELKE SR., A. BELTRÁN-PRZEKURAT, C.A. HIEMSTRA, J. LIN, AND T.E. NOBIS

Colorado State University
Department of Atmospheric Science
Fort Collins, CO 80523
pielke@atmos.colostate.edu
Phone: (970)491-8293/Fax: (970)491-3314

J. ADEGOKE

Laboratory for Climate Analysis & Modeling (LCAM), Department of Geosciences
420K Robert H. Flarsheim Hall, 5100 Rockhill Road
University of Missouri-Kansas City
Kansas City, MO 64110-2499

U.S. NAIR

Earth System Science Center
University of Alabama at Huntsville
320 Sparkman Dr
Huntsville, AL 35805

D. NIYOGI

Purdue University,
Department of Agronomy, and Department of Earth and Atmospheric Sciences,
Lilly Hall of Life Sciences, 915 W. State Street
Purdue University
West Lafayette, IN 47907-2054

Submitted to *The Hydrological Sciences Journal*, February 22, 2006

Abstract

This paper documents the diverse role of land-use/land-cover change on precipitation. Since land conversion continues at a rapid pace (e.g., see Table 1 in Pielke et al. 2006a), this type of human disturbance of the climate system will continue and become even more significant in the coming decades. The regional alteration of landscape also has global climate effects through teleconnections as concluded in NRC (2005); a conclusion which is bolstered by studies such as that of Chase et al. (2000) and Fedemma et al. (2005).

KEY WORDS: Land-atmosphere interactions; numerical modeling; rainfall/runoff; regional climate; land-use/land-cover change

INTRODUCTION

The role of landscape change in altering convective rainfall is well documented (e.g., Pielke 2001; Pitman 2003). This overview paper summarizes this topic by landscape conversion type focusing on how regional change results in changes in rainfall in the same area.

SHORT GRASS CONVERSION TO DRYLAND AGRICULTURE AND IRRIGATED AGRICULTURE

The conversion from native shortgrass grassland to cropland is manifested in biophysical effects that influence energy and water cycling. Seasonality of growing season, albedo, leaf area index, surface roughness, and moisture fluxes were altered with conversion to cropland. Cropland and grassland albedos are similar during the crop growing season (Oke 1987), but croplands have bare soil for much of the year and native vegetation has a higher albedo than bare soil (Bonan 2002). Dryland and irrigated crops are taller, and possess more leaf area than native shortgrass steppe (Paruelo et al. 2001). Moreover, surface roughness is higher with taller agricultural plants (Bonan 2002; Chase et al. 1999). Lastly, moisture fluxes are higher in the agricultural systems, especially irrigated croplands (Baron et al. 1998; Chase et al. 1999; Stohlgren et al. 1998).

Growing season dynamics change with conversion to agriculture. With shortgrass steppe featuring a mixture of cool- and warm-season grasses, photosynthesis occurs during the entire growing season and peak biomass occurs in early summer (Paruelo et al. 2001). In contrast, croplands have one dominant plant with dramatically different growing seasons and peak biomass, with dryland and irrigated crops peaking earlier and later than shortgrass, respectively (Paruelo et al. 2001).

The effect on air temperature and precipitation produced with shortgrass conversion varies with the conversion, spatially and temporally. The magnitude of change from shortgrass steppe to irrigated agriculture is much more dramatic than the shift to dryland

agriculture (Baron et al. 1998). At relatively fine scales during short periods of the growing season, lower temperatures and higher atmospheric moisture levels are closely tied to irrigated croplands (Chase et al. 1999; Segal et al. 1989). At larger scales, given the same brief temporal scale, this difference produces a regional cooling effect and precipitation increase on the adjacent Rocky Mountains (Stohlgren et al. 1998). In a coarse resolution (50 km grid increment) regional modeling comparison of natural and current vegetation change, Eastman et al. (2001) showed a 0-1°C increase in maximum temperatures over unconverted shortgrass steppe, while a 2-3°C increase occurred in areas converted to dryland crops. Precipitation changes were heterogeneous.

TALL GRASS CONVERSION TO DRYLAND AND IRRIGATED AGRICULTURE

The tall grass conversion is similar to the short grass conversion except the loss of aboveground biomass (leaf area index- LAI) is greater when the tall grass prairie was removed, and almost 100% of the original tall grass region is gone. Further studies on the role of grassland conversion include the initial evidence of the role of irrigation in modifying surface climate trends which came from observational studies (Marotz et al. 1975; Barnston and Schickendanz 1984; Alpert and Mandel 1986; Pielke and Zeng 1989). Barnston and Schickendanz (1984) found that irrigation increased precipitation in the Texas Panhandle when the synoptic condition provided low-level convergence and uplift such that the additional moisture produced by irrigation was allowed to ascend to cloud base.

Adegoke et al. (2006) found that midsummer 2 m temperature over Nebraska might be cooler by as much as 3.4°C under current irrigation (control run) conditions as contrasted with natural landscape conditions. Energy budget differences were identified between current and potential natural vegetation scenarios: the surface latent heat flux was 42% higher, and the water vapor flux (at 500 m) 38% greater in the control run compared to the natural landscape run. Important physical changes between the natural tallgrass prairie of this region

and the current land-use patterns include alterations in the surface albedo, roughness length, and soil moisture in the irrigated areas.

These changes are capable of generating complex changes in the lower atmosphere (PBL) energy budget. For example, the simulated increase in the portion of the total available energy partitioned into latent heat rather than sensible heat resulted directly from enhanced transpiration and soil evaporation in the control run. Although not examined in detail in this study, elevated dewpoint temperature and moisture fluxes within the PBL can increase convective available potential energy, promote atmospheric instability, and enhance daytime cloud cover (Alapaty et al. 1997; Stohlgren et al. 1998; Pielke 2001; Holt et al. 2006).

MID-LATITUDE DECIDUOUS FOREST CONVERSION TO AGRICULTURE

Asner et al. (2004) document that agricultural activities associated with grazing operations is an important driver for conversion from deciduous forest to grassland and pastures particularly under poor soil conditions. This conversion is surmised to have hydrological impacts by reducing the rate of spring snowmelt and cloud condensation levels and hence moisture availability, reduces moisture interception due to decreased LAI, increases runoff and soil evaporation, and decreases transpiration. This latter effect leads to higher soil moisture fluxes and discharge over the landscape, which causes erosion and poor soil conditions in the region. Typically, the change from forest to grasslands leads to a significant reduction in moisture flux, while a change to cropland can have variable influences on the regional moisture flux (because of crop photosynthetic pathways and transpiration rates, cropping patterns, irrigation, etc.).

Hogg et al. (2000) used field measurements over deciduous forest to demonstrate that the distinctive climate of interior western Canada is the feedback associated with the leaf phenology of the aspen forest. Latent heat fluxes are largest under high LAI conditions

leading to cooling, but with the resultant higher concentrations of water vapor, increased moisture availability and precipitation. Xue et al. (1996) indicate that the LAI changes of deciduous vegetation cause regional changes and propagate high uncertainty into general circulation model (GCM) simulations. Bounoua et al. (2000) concluded that for the midlatitudinal forest regions, the resulting impact of LAI changes was a decreased albedo, cooling of about 1.8 K during the growing season, and slight warming during winter due to snow albedo masking, decreased effective precipitation, and an increase in the low-frequency variability of weather in the northern latitudes. Baidya Roy et al. (2002) simulated a 300-year (1700-1990) time series of U.S. land use/land cover and found that changing land-use/land-cover patterns can lead to several degrees of warming or cooling at the surface accompanied by significant changes in precipitation patterns. Satellite products can detect this change following techniques similar to those discussed in Shepherd (2005) and Cai and Kalnay (2005) using blended reanalysis datasets.

TROPICAL EVERGREEN FOREST CONVERSION TO AGRICULTURE

Tropical forests, occupying approximately 800 million hectares, are being cleared at the rate of approximately 14 million hectares per year. Observational studies, spanning several decades, and numerical modeling studies both show that tropical deforestation influence cloud formation and rainfall (Sud and Smith 1985; Meher-Homji 1991; Cutrim et al. 1995; McGuffie et al. 1995; Costa and Foley 2000; Lawton et al. 2001; Pielke 2001; Silva Dias et al. 2002; Durieux et al. 2003; Sen et al. 2004; Fisch et al. 2004; Ray et al. 2006). Observational studies include Meher-Homji (1991), Pielke (2001), Durieux et al. (2003) and Ray et al. (2006). These studies report a wide range of changes in rainfall associated with deforestation (1-20% decrease), as well as the alteration of seasonality and frequency of convection. Regional-scale modeling results show that the eastern Asian summer monsoon is sensitive to deforestation in the Indochina region (Sen et al. 2004). Heterogeneity-induced

mesoscale circulations have the potential to modify cloudiness (Souza et al. 2000; Pielke 2001; Silva Dias et al. 2002; Baidya Roy and Avissar 2002; Werth and Avissar 2002). Mesoscale numerical modeling experiments show that lowland deforestation and associated increases (decreases) in the Bowen ratio lead to elevation (lowering) of the orographic clouds forced by terrain downwind (Lawton et al. 2001; van der Molen 2002; Nair et al. 2003; Bruijnzeel 2004; Ray et al. 2006), leading to changes in the direct harvesting of cloud water by montane vegetation.

There is some indication that deforestation in a continent surrounded by oceans has more potential to impact tropical circulation compared to deforestation that is further removed from ocean sources of water vapor (van der Molen et al. 2006), and this needs to be validated through the use of cloud-resolving and regional modeling experiments.

BOREAL FOREST CONVERSION DUE TO FIRE

Fire in the boreal forest is an integral part of this ecosystem (Stocks 1991). As shown in Vidale et al. (1997), mesoscale circulations develop in response to the spatial variations of surface sensible heat fluxes that occur due to these fires and other landscape pattern disturbances, which also include the ubiquitous lakes that exist in the boreal forests. Knowles (1993) reported that cumulus clouds form preferentially downwind of recent burn areas within a boreal forest landscape. These burn areas have a lower albedo than the surrounding landscape. Such a preference also indicates that subsequent lightning strikes from deep cumulus clouds will more likely initiate fires immediately downstream of a recent burn scar, rather than elsewhere in the forest. After a period of time, as the forest regenerates, the burned area may be lighter than the surrounding unburnt landscape, as aspen and other second growth forest and shrubs grow.

The environment of the boreal forest, particularly in the spring, is very conducive to fires as the roots are embedded in cold, or even frozen soils, yet the air temperature and solar insolation at this time of year is high. Almost all of the net radiation received at the surface is transferred back into the atmosphere as sensible, rather than latent heat flux (Sellers et al. 1995). Pielke and Vidale (1995) show that one consequence of the resultant large heating of the atmosphere by the sensible heating is a particularly deep planetary boundary layer, as well as a preference for the summer polar front to often situate along the boreal-tundra ecotone boundary. The region south of the polar front in the spring (and also in the summer) is a weather location where thunderstorms occur. The spatial variations of the landscape will provide focused regions for thunderstorm development that would otherwise not occur.

URBANIZATION

While the fraction of the Earth's surface currently classified as urban accounts for less than 2% of available land surface, over 45% of the Earth's population is concentrated there (Arnfield 2003). Furthermore, future projections indicate this percentage will soon exceed more than 50% (Cohen 2003). Observational studies over the past three decades have demonstrated that urban areas radically restructure the local energy budget and thus lead to different boundary layer structure (Shepherd 2005; Arnfield 2003; Oke 1988). The anthropogenic influence also includes altering the aerosol environment. These changes likely lead to alterations in urban precipitation frequency, intensity, and patterns. Shepherd (2005) presents the most recent and complete review of urban precipitation issues.

Observational studies of urban precipitation stretches over three decades (e.g., Huff and Changnon 1973). Studies in the U.S. have been mainly concentrated in three urban areas; St. Louis, Atlanta, and Houston and generally demonstrated increases in rainfall over and downwind of urban areas attributed largely to the Urban Heat Island (UHI) initiated convergence zone and to a lesser extent, increased surface roughness (Shepherd and Burian

2003; Dixon and Mote 2003; Bornstein and Lin 2000; Huff 1986; Huff and Changnon 1973). Evidence also suggests an increase in heavy rain events (Huff 1986). In addition, the UHI has been found to decrease the likelihood of freezing rain events in urban areas (Changnon 2003). Bornstein and Lin (2000) also found urban influence on established thunderstorms approaching an urban area. Model studies have demonstrated significant changes to the convective boundary layer structure (e.g., Hildebrand and Ackerman 1984; Baik et al. 2001) and even influences on the behavior of cold fronts (Gaffen and Bornstein 1988).

A big unknown in the urban-induced precipitation problem is the role of urban aerosols. Studies like van den Heever and Cotton (2005) have begun to examine the relative sensitivities involved, but as shown in Shepherd (2005), while evidence strongly points to a link between aerosols and urban precipitation modification, the details of the connection remain highly uncertain.

TROPICAL FOREST FIRES AND RESULTANT BIOMASS BURNING EFFECT ON RAINFALL

Biomass burning produces smoke plumes with large quantities of aerosols (tiny particles) which can potentially affect the regional, even global hydrological cycle (Ramanathan et al. 2001). Aerosols serve as cloud condensation nuclei which effect the formation of cloud droplets (Cotton and Anthes 1992); thus the extensive input of aerosols from fires could significantly affect cloud properties and rainfall. In addition to affecting cloud microphysics, aerosols absorb and scatter radiation. Increased aerosols in the atmosphere from forest fires reduce the radiative energy reaching the Earth's surface (Ramanathan et al. 2001). Because surface radiative input drives evaporation, which is balanced with precipitation on a global scale, aerosol-induced reductions in surface radiation are likely to decrease global rainfall (Lohmann and Feichter 2005).

Andreae et al. (2004) analyzed Amazonian aircraft observations and found delays in the onset of precipitation within smoky clouds. Khain et al. (2005) show that while elevated aerosol concentrations initially lead to lowered precipitation efficiency due to warm-rain suppression, the delay in raindrop formation decreases the drag on updrafts by falling raindrops and increases the latent heat release by the additional water that reaches higher altitudes where freezing takes place.

Lin et al. (2006), based on satellite observations during the entire Amazonian biomass burning season, empirically confirmed the aforementioned results of Andreae et al. (2004) and Khain et al. (2005). Increased aerosols from fires were correlated with increased observed total (warm+ice-phase) rainfall from the TRMM-TMI sensor, even after accounting for the atmospheric stability environment through the cloud work function. Changes in cloud properties were also correlated with aerosol loading; i.e., higher cloud tops, increased presence of ice, and enhanced cloud cover.

AFFORESTATION AND REFORESTATION

Afforestation and reforestation (A&R) are proposed as possible tools to mitigate desertification (FAO 1989) and to reduce atmospheric concentrations of CO₂ by sequestering carbon in forest biomass (UNDP 2003). In southern South America, A&R plans started in Chile, Uruguay, and Argentina in the last two decades, supported through government economic incentives or subsidies (World Bank 2000). In southwestern Australia, A&R are also seen as ways to ameliorate salinity (Walker et al. 2002).

Overall, conversion from grasslands or croplands to forest leads to a decrease in albedo and increases of LAI, roughness length, and rooting depth (Sellers 1992; Pitman 2003; Jackson et al. 1996). Changes in these parameters can modify the near-surface energy fluxes, which can influence temperature and humidity (Pielke 2001). In general, observations and modeling studies agree that A&R would decrease near-surface temperature and increase

latent heat (e.g., Beltrán 2005; Fahey and Jackson 1997; Noretto et al. 2005; Xue and Shukla 1996). Simulated impacts on precipitation are not so clear, and depend on geographical location, regional atmospheric characteristics, extent of the afforested-reforested area (Pitman and Narisma 2005; Xue and Shukla 1996) and biophysical parameters of the land-use/land-cover change (Xue et al. 1996).

Observational (Jackson et al. 2005) and several modeling studies (Beltrán 2005; Xue et al. 1996) demonstrated that tree plantation establishment may affect the hydrological cycle. Precipitation processes depend on local, regional, and large-scale atmospheric characteristics and therefore regional atmospheric modelling represent an important tool to study the impacts of realistic patterns of A&R on precipitation. Moreover, model domain size, grid spacing, and parameterizations (e.g., convective schemes) may largely influence simulated precipitation (e.g., Castro et al. 2005). Thus, more comprehensive regional modeling sensitivity studies are needed to assess those impacts.

Acknowledgements

The work reported in this article was sponsored by NASA Grant NNG04GL61G, NNG04G184G, and NAG5-11370, USGS Grant 04CRAG0032, and NSF Grants DEB 0217631 and ATM-0233780. Dr. John C. Lin was supported by the NOAA Postdoctoral Program in Climate and Global Change, administered by the University Corporation for Atmospheric Research. Dallas Staley completed the editorial work on this paper with her customary outstanding professionalism and efficiency.

REFERENCES

Adegoke JO, Pielke RA Sr., Carleton AM (2006) Observational and modeling studies of the impacts of agriculture-related land use change on climate in the central U.S. *Agric. Forest Meteor.* Special Issue, submitted.

- Alapaty K, Pleim JE, Raman S, Niyogi DS, Byun DW (1997) Simulation of atmospheric boundary layer processes using local- and nonlocal-closure schemes. *J Appl Meteor* **36**:214-233
- Alpert P, Mandel M (1986) Wind variability – An indicator for mesoclimatic change in Israel. *J Climate Appl Meteor* **25**:1568-1576
- Andreae MO, Rosenfeld D, Artaxo P, Costa AA, Frank GP, Longo KM, Silva-Dias MAF (2004) Smoking rain clouds over the Amazon. *Science* **303**:1337-1342
- Arnfield AJ (2003) Two decades of urban climate research: A review of turbulence, exchanges of energy and water, and the Urban heat island. *Intl J Climate* **23**:1-26
- Asner GP, Elmore AJ, Olander LP, Martin RE, AT (2004) Grazing systems, ecosystem responses, and global change. *Ann Rev Environ Resources* **29**:261-299
- Baidya Roy S, Avissar R (2002) Impact of land use/land cover change on regional hydrometeorology in Amazonia. *J Geophys Res* **107** (D20): doi:10.1029/2000JD000266.
- Baidya Roy S, Weaver CP, Pacala SW, Hurtt G (2002) Impact of historical land-use/land-cover change on US summer climate. Amer Geophys Union, AGUSM.B41A.12B
- Baik JJ, Kim YH, Chun HY (2001) Dry and moist convection forced by an urban heat island. *J Applied Meteor* **40**:1462-1475
- Barnston AG, Schickedanz PT (1984) The effect of irrigation on warm season precipitation in the southern Great Plains. *J Climate Appl Meteor* **23**:865-888
- Baron JS, Hartman MD, Kittel TGF, Band LE, Ojima DS, Lammers RB (1998) Effects of land cover, water redistribution, and temperature on ecosystem processes in the South Platte Basin. *Ecological Applications* **8**:1037-1051
- Beltrán A (2005) Using a coupled atmospheric-biospheric modeling system (GERRAMS) to model the effects of land-use/land-cover changes on the near-surface atmosphere. PhD, Department of Atmospheric Science, Colorado State University, 186 pp.

- Bonan GB (2002) *Ecological Climatology: Concepts and Applications*. Cambridge University Press, 678 pp.
- Bornstein R, Lin Q (2000) Urban heat islands and summertime convective thunderstorms in Atlanta: Three case studies. *Atmos Environ*, **34**:507-516
- Bounoua L, Collatz GJ, Los SO, Sellers PJ, Dazlich DA, Tucker CJ, Randall DA (2000) Sensitivity of climate to changes in NDVI. *J Climate* **13**:2277-2292
- Bruijnzeel LA (2004) Hydrological functions of tropical forests: Not seeing the soil for the trees? *Agriculture Ecosystems and Environment*. 104:185-228
- Cai, M, Kalnay E (2005) Can reanalysis have anthropogenic climate trends without model forcing? *J Climate*, 18:1844–1849
- Castro CL, Pielke RA Sr, Leoncini G (2005) Dynamical downscaling: assessment of value retained and added using the Regional Atmospheric Modeling System (RAMS). *J Geophys Res* **110**:D05108, doi:10.1029/2004JDD004721
- Changnon SA (2003) Urban modification of freezing-rain events. *J Appl Meteor* **42**:863-870
- Chase TN, Pielke RA, Kittel TGF, Baron JS, Stohlgren TJ (1999) Potential impacts on Colorado Rocky Mountain weather due to land use changes on the adjacent Great Plains. *J Geophys Res-Atmos* **104**:16673-16690
- Chase TN, Pielke RA, Kittel TGF, Nemani RR, and Running SW (2000) Simulated impacts of historical land cover changes on global climate in northern winter. *Climate Dynamics* **16**:93-105
- Cohen JE (2003) Human population: The next half century. *Science* **302**:1172-1175
- Costa MH, Foley JA (2000) Combined effects of deforestation and doubled atmospheric CO₂ concentrations on the climate of Amazonia. *J Climate* **13**:18-34
- Cotton WR, Anthes RA (1992) *Storm and Cloud Dynamics*, Acad. Press, San Diego, 883 pp.

- Cutrim ED, Martin W, Rabin R (1995) Enhancement of cumulus clouds over deforested lands in Amazonia. *Bull Amer Meteor Soc* **76**:1801-1805
- Dixon PG, Mote TL (2003) Patterns and causes of Atlanta's urban heat island-initiated precipitation. *J Applied Meteor* **42**:1273-1284
- Duriex L, Machado L, Laurent H (2003) The impact of deforestation on cloud cover over the Amazon arc of deforestation. *Remote Sensing of Environment* **86**:132-140
- Eastman JL, Coughenour MB, Pielke RA (2001) The effects of CO₂ and landscape change using a coupled plant and meteorological model. *Global Change Biology* **7**:797-815
- Fahey B, Jackson RB (1997) Hydrological impacts of converting native forests and grasslands to pine plantations, South Island, New Zealand. *Agr For Meteor* **84**:69-82
- FAO (Food and Agriculture Organization of the United Nations) (1989) Role of forestry in combating desertification. *FAO Conservation Guide*, **21**, 331 pp.
- Feddema JJ, Oleson KW, Bonan GB, Mearns LO, Buja LE, Meehl GA, Washington WM (2005) The importance of land-cover change in simulating future climates. *Science* **310**:1674-1678
- Fisch G, Tota J, Machado LAT, Silva Dias MAF, da F Lyra RF, C.A Nobre CA, Dolman AJ, Gash JHC (2004) The convective boundary layer over pasture and forest in Amazonia. *Theor Appl Climatol* **78**:47-59
- Gaffen D, Bornstein R (1988) Case study of urban interactions with a synoptic scale cold front. *Meteor Atmos Phys* **38**:185-194
- Hildebrand PH, Ackerman B (1984) Urban effects on the convective boundary layer. *J Atmos Sci* **41**:76-91
- Hogg EH, Price DT, Black TA (2000) Postulated feedbacks of deciduous forest phenology on seasonal climate patterns in the western Canadian interior. *J Climate* **13**:4229-4243

- Holt T, Niyogi DS, Chen F, Manning K, LeMone M, Qureshi A (2006) Effect of land-atmosphere interactions on the IHOP 24-25 May 2002 convection case. *Mon Wea Rev*, **134**:113–133
- Huff TE (1986) Urban hydrometeorology review. *Bull Amer Meteor Soc* **67**:703-711
- Huff FA Changnon SA Jr (1973) Precipitation modification by major urban areas. *Bull Amer Meteor Soc* **54**:1220-1232
- Jackson RB, Canadell J, Ehleringer JR, Mooney HA, Sala OE, Schulze ED (1996) A global analysis of root distributions for terrestrial biomes. *Oecologia* **108**:389-411
- Jackson RB, Jobbágy EG, Avissar R, Baidya Roy S, Barrett DJ, Cook CW, Farley KA, le Maitre DC, McCarl BA, Murray BC (2005) Trading water for carbon with biological carbon sequestration. *Science* **310**:1944-1947
- Khain A, Rosenfeld D, Pokrovsky A (2005) Aerosol impact on the dynamics and microphysics of deep convective clouds. *Quart J Roy Meteor Soc* **131**:1-25
- Knowles J.B (1993) M.S. Thesis: The influence of forest fire induced albedo differences on the generation of mesoscale circulations. Dept Atmos Sci, CO State Univ, 86 pp.
- Lawton RO, Nair US, Pielke RA, Welch RM (2001) Climatic impact of tropical lowland deforestation on nearby montane cloud forests. *Science* **294**:584-587
- Lin JC, Matsui T, Pielke RA Sr, Kummerow C (2006) Effects of biomass burning-derived aerosols on precipitation and clouds in the Amazon Basin: a satellite-based empirical study. *J Geophys Res*, submitted.
- Lohmann U, Feichter J (2005) Global indirect aerosol effects: A review. *Atmos Chem Phys* **5**:715-737
- Marotz GA, Clark J, Henry J, Standfast R (1975) Cloud fields over irrigated areas in southwestern Kansas – Data and speculations. *Prof Geogr* **27**:457-461

- McGuffie K, Henderson-Sellers A, Zhang H, Durbidge TB, Pitman AJ (1995) Global climate sensitivity to tropical deforestation. *Global and Planetary Change* **10**:97-128
- Meher-Homji VM (1991) Probable impact of deforestation on hydrological processes. *Clim Change* **19**:163-173
- Molders N, Olson MA (2004) Impact of urban effects on precipitation in high latitudes. *J Hydrometeor* **5**:409-429
- Nair US, Lawton RO, Welch RM, Pielke RA Sr (2003) Impact of land use on Costa Rican tropical montane cloud forests: Sensitivity of cumulus cloud field characteristics to lowland deforestation. *J Geophys Res* **108(D7)**:4206, doi:10.1029/2001JD001135
- National Research Council (2005) *Radiative forcing of climate change: Expanding the concept and addressing uncertainties*, National Academies Press, Washington, DC
- Nosetto MD, Jobbagy EG, Paruelo JM (2005) Land-use change and water losses: the case of grassland afforestation across a soil textural gradient in central Argentina. *Global Change Biol* **11**:1101-1117
- Oke TR (1987) *Boundary Layer Climates*. 2nd Ed. Methuen, xxiv, 435 pp.
- Oke TR (1988) The urban energy balance. *Prog Phys Geogr* **12**:471-508
- Paruelo JM, Burke IC, Lauenroth WK (2001) Land-use impact on ecosystem functioning in eastern Colorado, USA. *Global Change Biology* **7**:631-639
- Pielke Sr RA (2001) Influence of the spatial distribution of vegetation and soils on the prediction of cumulus convective rainfall. *Rev Geophys* **39**:151-177
- Pielke Sr RA, Zeng X (1989) Influence on severe storm development of irrigated land. *National Weather Digest* **14**:16-17
- Pielke RA, Vidale PL (1995) The boreal forest and the polar front. *J Geophys Res* **100**:25755-25758

- Pielke Sr RA, Adegoke JO, Beltrán-Prezekurat A, Hiemstra CA, Lin J, Nair US, Niyogi D, Nobis TE (2006a) An overview of regional land use and land cover impacts on rainfall. *Tellus B*, submitted.
- Pielke Sr RA, Adegoke JO, Chase TN, Marshall CH, Matsui T, Niyogi D (2006b) A new paradigm for assessing the role of agriculture in the climate system and in climate change. *Agric. Forest Meteor.*, Special Issue (in press)
- Pitman AJ (2003) The evolution of, and revolution in, land surface schemes designed for climate models. *Int J Climatol* **23**:479-510
- Pitman AJ, Narisma GT (2005) The role of land surface processes in regional climate change: a case study of future land cover change over south western Australia. *Meteor Atmos Phys* **89**:235-249
- Ramanathan V, Crutzen PJ, Kiehl JT, Rosenfeld D (2001) Aerosols, climate, and the hydrological cycle. *Science* **294**:2119-2124
- Ray DK, Nair US, Lawton RO, Welch RM, Pielke RA Sr (2006) Impact of land use on Costa Rican tropical montane cloud forests: Sensitivity of orographic cloud formation to deforestation in the plains. *J Geophys Res* 111:D02108, doi:10.1029/2005JD006096.
- Segal M, Schreiber WE, Kallos G, Garratt JR, Rodi A, Weaver J, Pielke RA (1989) The impact of crop areas in northeast Colorado on midsummer mesoscale thermal circulations. *Mon Wea Rev* **117**:809-825
- Sellers PJ (1992) Biophysical models of land surface processes. In: *Climate System Modeling*, Trenberth KE (ed), Cambridge University Press, 451-490 pp
- Sellers P, Hall F, Ranson KJ, Margolis H, Kelly B, Baldocchi D, den Hartog G, Cihlar J, Ryan MG, Goodison B, Crill P, Lettenmaier D, Wickland DE (1995) The Boreal Ecosystem–Atmosphere Study (BOREAS): An overview and early results from the 1994 field year. *Bull Amer Meteor Soc* **76**:1549–1577

- Sen OL, Wang Y, Wang B (2004) Impact of Indochina deforestation on the East Asian summer monsoon. *J Climate* **17**:1366-1380
- Shepherd JM (2005) A review of current investigations of urban-induced rainfall and recommendations for the future. *Earth Interactions* **9**: paper 12, 27 pp.
- Shepherd JM, Burian SJ (2003) Detection of urban-induced rainfall anomalies in a major coastal city. *Earth Interactions* **7**: paper 4, 17 pp.
- Silva Dias M, Rutledge S, Kabat P, Silva Dias PL, Nobre C, Fisch G, Dolman AJ, Zipser E, Garstang M, Manzi AO, Fuentes J, Andreae MO, Artaxo P, Gielow R, Gatti L (2002) Cloud and Rain Processes in a Biosphere-Atmosphere Interaction Context in the Amazon Region. *J Geophys Res* **29**:10.1029/2000JD000335
- Souza EP, Rennó NO, Silva Dias MAF (2000) Convective circulations induced by surface heterogeneities. *J Atmos Sci* **57**:2915–2922
- Stocks BJ (1991) The extent and impact of forest fires in northern circumpolar countries. In *Global biomass burning-atmospheric, climatic, and biospheric implications*. MIT Press, 197-202
- Stohlgren TJ, Chase TN, Pielke RA, Kittel TGF, Baron JS (1998) Evidence that local land use practices influence regional climate, vegetation, and stream flow patterns in adjacent natural areas. *Global Change Biology* **4**:495-504
- Sud YC, Smith WE (1985) Influence of local land surface processes on the Indian monsoon - A numerical study. *J Climate Appl Meteor* **24**(10):1015-1036
- UNDP [United Nations Development Programme] (2003) The Clean Development Mechanism: A User's Guide. Bureau for Development Policy. NY 20 pp
- Van den Heever SC, Cotton WR (2005) Modeling the impacts of urban aerosol on convection and precipitation. Presentation, 2nd Urban Symposium, San Diego, CA, January 2005.

- Van der Molen MK (2002) Meteorological impacts of land use change in maritime tropics, PhD, Vrije Universiteit, Amsterdam, The Netherlands, 262 pp.
- Van der Molen MK, Dolman AJ, Waterloo MJ, Bruijnzeel LA (2006) Climate is affected more by maritime than by continental land use change: A multiple scale analysis. *Global and Planetary Change*, in press
- Vidale PL, Pielke RA, Barr A, Steyaert LT (1997) Case study modeling of turbulent and mesoscale fluxes over the BOREAS region. *J Geophys Res* **102**:29167-29188
- Walker GR, Zhang L, Ellis TW, Jatton TH, Petheram C (2002) Estimating impacts of changed land use on recharge: review of modeling and other approaches appropriate for management of dryland salinity. *Hydrogeol J* **10**:68-90
- World Bank (2000) Policy Elements for Sustainable Development in the XXI Century. Argentina Water Resources Management. February 18, 2000. Latin America and the Caribbean Regional Argentina Country Department and Finance, Private Sector and Infrastructure, and Environment and Socially Sustainable Development.
- Xue Y, Shukla J (1996) The influence of land surface properties on Sahel climate. Part II: Afforestation. *J Climate* **9**:3260-3275
- Xue Y, Fennessy MJ, Sellers PJ (1996) Impact of vegetation properties on U.S. summer weather prediction. *J Geophys Res* **101**:D3, 7419–7430